

TWIGAMYCIN

POM

Oxytetracycline 10% Injection in Cattle, Sheep, Goats & Pigs

FOR VETERINARY USE ONLY

DESCRIPTION

Oxytetracycline belongs to the group of tetracyclines and acts bacteriostatic against many Gram-positive and Gram-negative bacteria like *Bordetella*, *Campylobacter*, *Chlamydia*, *E. coli*, *Haemophilus*, *Mycoplasma*, *Pasteurella*, *Rickettsia*, *Salmonella*, *Staphylococcus* and *Streptococcus* spp. The action of oxytetracycline is based on inhibition of bacterial protein synthesis. Oxytetracycline is mainly excreted in urine, for a small part in bile and in lactating animals in milk.

COMPOSITION

Contains per ml.:

Oxytetracycline base 100 mg.

Solvents ad. 1 ml.

INDICATIONS

Arthritis, gastrointestinal and respiratory infections caused by oxytetracycline sensitive micro-organisms, like *Bordetella*, *Campylobacter*, *Chlamydia*, *E. coli*, *Haemophilus*, *Mycoplasma*, *Pasteurella*, *Rickettsia*, *Salmonella*, *Staphylococcus* and *Streptococcus* spp., in calves, cattle, goats, sheep and swine.

CONTRA-INDICATIONS

- Hypersensitivity to tetracyclines.
- Administration to animals with a serious impaired renal and/or liver function.
- Concurrent administration with penicillins, cephalosporines, quinolones and cycloserine.
- Do not use in cats, dogs and horses.

SIDE EFFECTS

- After intramuscular administration local reactions can occur, which disappear in a few days.
- Discoloration of teeth in young animals.
- Hypersensitivity reactions.

DOSAGE AND ADMINISTRATION

For intramuscular or subcutaneous administration:

Full-grown animals : 1 ml. per 10 - 20 kg. body weight for 3 - 5 days.

Young animals : 2 ml. per 10 - 20 kg. body weight for 3 - 5 days.

Do not administer more than 20 ml. in cattle, more than 10 ml. in swine and more than 5 ml. in calves, goats and sheep per injection site.

WITHDRAWAL TIMES

- For meat : 15 days.

- For milk : 5 days.

STORAGE

Store in a cool dry place away from direct sunlight

Store Below 30°C.

Protect from Light.

WARNING

Keep out of reach of children.

Avoid the introduction of contamination during use.

PACKING

Vial of 50ml and 100ml.

A product of:

Twiga Chemical Industries Ltd.

P.O. Box 30172 -00100, NAIROBI, KENYA

Tel: +254 20 3942000, Fax: +254 20 3942405

